

Medication Facts

Active Ingredients Purposes

(In Each Softgel)

feniramine 10 mg	Antihistamine
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant
Tromethaphan 35mg	Cough Suppressant

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease
- excessive phlegm (mucous)
- glaucoma
- high blood pressure
- thyroid disease
- diabetes
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur
- avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
- stomach pain occurs
- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days

Ask a doctor. These may be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Medication Facts

Active Ingredients (In Each Softgel)	Purposes
Ilfenesine 10 mg	Antihistamine
Pseudoephedrine	Nasal Decongestant
Hydrochloride 40 mg	
Tromethaphan 35mg	Cough Suppressant

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Warnings

- Do Not Use:**
- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
 - for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
 - during the first 4 months of pregnancy

Ask A Doctor Before Use

- If You Have:**
- heart disease
 - excessive phlegm (mucous)

- glaucoma
- diabetes
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur
- avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
- stomach pain occurs
- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days

Ask a doctor. These may be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away. Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Medication Facts

Active Ingredients (In Each Softgel)	Purposes
Pheniramine 10 mg	Antihistamine
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant
Triprolidine 35mg	Cough Suppressant

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease
- excessive phlegm (mucous)
- glaucoma
- high blood pressure
- diabetes
- thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur
- avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
 - stomach pain occurs
 - nervousness, dizziness or sleeplessness occur
 - symptoms do not improve within 7 days
- Ask a doctor. These may be signs of a serious condition.**

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

<h2>Medication Facts</h2>	
Active Ingredients (In Each Softgel) Iifenesine 10 mg Antihistamine Pseudoephedrine Hydrochloride 40 mg Nasal Decongestant Tromethphan 35mg Cough Suppressant	Purposes ■ glaucoma ■ high blood pressure ■ diabetes ■ thyroid disease ■ a cough that lasts from smoking, asthma or emphysema ■ cough with fever, rash or headache that lasts ■ a breathing problem such as emphysema or chronic bronchitis ■ difficulty in urination due to prostate gland enlargement
Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion	
Directions Do not use more than directed.	
Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product
Warnings Do Not Use: ■ if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease) ■ for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI. ■ during the first 4 months of pregnancy	
Ask A Doctor Before Use If You Have: ■ heart disease ■ excessive phlegm (mucous)	
When Using This Product: ■ drowsiness may occur ■ avoid alcohol ■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect ■ use caution when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children	
Stop Using This Product If: ■ cough is accompanied by fever, rash or headache that lasts ■ stomach pain occurs ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not improve within 7 days Ask a doctor. These may be signs of a serious condition. If pregnant or breast feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help right away. Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).	

Active Ingredients (In Each Softgel) Iifenessine 10 mg Antihistamine Pseudoephedrine Nasal Decongestant Hydrochloride 40 mg Tromethphan 35mg Cough Suppressant	Purposes 	If You Are: ■ taking any drugs for asthma ■ taking sedatives or tranquilizers						
Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion	When Using This Product: ■ drowsiness may occur ■ avoid alcohol ■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect ■ use caution when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children	Stop Using This Product If: ■ cough is accompanied by fever, rash or headache that lasts ■ stomach pain occurs ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not improve within 7 days Ask a doctor. These may be signs of a serious condition. If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help right away.						
Do Not Use: ■ if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease) ■ for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI. ■ during the first 4 months of pregnancy	Warnings							
Ask A Doctor Before Use If You Have: ■ heart disease ■ excessive phlegm (mucous) ■ glaucoma ■ high blood pressure ■ diabetes ■ thyroid disease ■ a cough that lasts from smoking, asthma or emphysema ■ cough with fever, rash or headache that lasts ■ a breathing problem such as emphysema or chronic bronchitis ■ difficulty in urination due to prostate gland enlargement	Directions Do not use more than directed. <table> <tr> <td>Adults and children over 12 years of age</td> <td>Take 2 softgels every 4 hours</td> </tr> <tr> <td>Children 6 to under 12 years</td> <td>Take 1 softgel every 4 hours</td> </tr> <tr> <td>Children under 6 years of age</td> <td>Should not use this product</td> </tr> </table>	Adults and children over 12 years of age	Take 2 softgels every 4 hours	Children 6 to under 12 years	Take 1 softgel every 4 hours	Children under 6 years of age	Should not use this product	
Adults and children over 12 years of age	Take 2 softgels every 4 hours							
Children 6 to under 12 years	Take 1 softgel every 4 hours							
Children under 6 years of age	Should not use this product							
Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).								

Active Ingredients (In Each Softgel)		Purposes
Ilfenesine 10 mg	Antihistamine
Pseudoephedrine	Nasal Decongestant
Hydrochloride 40 mg		
Tromethaphan 35mg	Cough Suppressant
Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion		
Directions		
Do not use more than directed.		
Adults and children over 12 years of age	Take 2 softgels every 4 hours	
Children 6 to under 12 years	Take 1 softgel every 4 hours	
Children under 6 years of age	Should not use this product	
Warnings		
Do Not Use:		
■ if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)		
■ for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.		
■ during the first 4 months of pregnancy		
Ask A Doctor Before Use		
If You Have:		
■ heart disease ■ excessive phlegm (mucous)		
■ glaucoma ■ high blood pressure		
■ diabetes ■ thyroid disease		
■ a cough that lasts from smoking, asthma or emphysema		
■ cough with fever, rash or headache that lasts		
■ a breathing problem such as emphysema or chronic bronchitis		
■ difficulty in urination due to prostate gland enlargement		
If You Are:		
■ taking any drugs for asthma		
■ taking sedatives or tranquilizers		
When Using This Product:		
■ drowsiness may occur ■ avoid alcohol		
■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect		
■ use caution when driving a motor vehicle or operating machinery		
■ excitability may occur, especially in children		
Stop Using This Product If:		
■ cough is accompanied by fever, rash or headache that lasts		
■ stomach pain occurs		
■ nervousness, dizziness or sleeplessness occur		
■ symptoms do not improve within 7 days		
Ask a doctor. These may be signs of a serious condition.		
If pregnant or breast feeding, ask a health professional before use.		
Keep out of reach of children.		
In case of overdose, get medical help right away.		
Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).		

Active Ingredients (In Each Softgel)	Purposes
Itenesine 10 mg	Antihistamine
Pseudoephedrine	Nasal Decongestant
Hydrochloride 40 mg	
Tromethaphan 35mg	Cough Suppressant

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease
- excessive phlegm (mucous)
- glaucoma
- high blood pressure
- diabetes
- thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur
- avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
- stomach pain occurs
- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days

Ask a doctor. These may be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Active Ingredients Purposes		<ul style="list-style-type: none">■ diabetes ■ thyroid disease■ cough with fever, rash or headache that lasts■ a cough that lasts from smoking, asthma or emphysema■ a breathing problem such as emphysema or chronic bronchitis■ difficulty in urination due to prostate gland enlargement
(In Each Softgel)		
Iphenesine 10 mg Antihistamine		
Pseudoephedrine Nasal Decongestant		
Hydrochloride 40 mg		<ul style="list-style-type: none">■ If You Are:■ taking any drugs for asthma■ taking sedatives or tranquilizers
Tromethaphan 35mg Cough Suppressant		
Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion		
Directions		
Do not use more than directed.		
Adults and children over 12 years of age	Take 2 softgels every 4 hours	When Using This Product: <ul style="list-style-type: none">■ drowsiness may occur ■ avoid alcohol■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect■ use caution when driving a motor vehicle or operating machinery■ excitability may occur, especially in children
Children 6 to under 12 years	Take 1 softgel every 4 hours	
Children under 6 years of age	Should not use this product	
Warnings		
Do Not Use: <ul style="list-style-type: none">■ if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)■ for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.■ during the first 4 months of pregnancy		
Ask A Doctor Before Use If You Have: <ul style="list-style-type: none">■ heart disease ■ excessive phlegm (mucous)■ glaucoma ■ high blood pressure		
Stop Using This Product If: <ul style="list-style-type: none">■ cough is accompanied by fever, rash or headache that lasts■ stomach pain occurs■ nervousness, dizziness or sleeplessness occur■ symptoms do not improve within 7 days		
Ask a doctor. These may be signs of a serious condition.		
If pregnant or breast feeding, ask a health professional before use.		
Keep out of reach of children. In case of overdose, get medical help right away.		
Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).		

Medication Facts

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease ■ excessive phlegm (mucous)
- glaucoma ■ high blood pressure
- diabetes ■ thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur ■ avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect

- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
 - stomach pain occurs
 - nervousness, dizziness or sleeplessness occur
 - symptoms do not improve within 7 days
- Ask a doctor. These may be signs of a serious condition.**

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Active Ingredients (In Each Softgel)

Ifenazine 10 mg Antihistamine
Pseudoephedrine Nasal Decongestant
Hydrochloride 40 mg

Tromethaphan 35mg Cough Suppressant

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Medication Facts

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Do Not Use: Warnings

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

- If You Have:**
- heart disease ■ excessive phlegm (mucous)
 - glaucoma ■ high blood pressure
 - thyroid disease ■ diabetes
 - a cough that lasts from smoking, asthma or emphysema
 - cough with fever, rash or headache that lasts
 - a breathing problem such as emphysema or chronic bronchitis

- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur ■ avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
 - stomach pain occurs
 - nervousness, dizziness or sleeplessness occur
 - symptoms do not improve within 7 days
- Ask a doctor. These may be signs of a serious condition.**

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Active Ingredients	Purposes
(In Each Softgel)	
fenesine 10 mg	Antihistamine
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant
Tromethaphan 35mg	Cough Suppressant

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Medication Facts

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Do Not Use: Warnings

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease ■ excessive phlegm (mucous)
- glaucoma ■ high blood pressure
- diabetes ■ thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur ■ avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect

- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
- stomach pain occurs
- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days

Ask a doctor. These may be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Active Ingredients (In Each Softgel)

Ilfenesine 10 mg Antihistamine
Pseudoephedrine Nasal Decongestant
Hydrochloride 40 mg
Tromethaphan 35mg Cough Suppressant

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

<h2>Medication Facts</h2> <p>Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion</p>		<ul style="list-style-type: none"> ■ difficulty in urination due to prostate gland enlargement <p>If You Are:</p> <ul style="list-style-type: none"> ■ taking any drugs for asthma ■ taking sedatives or tranquilizers 								
<p>Directions</p> <p>Do not use more than directed.</p> <table border="1"> <tr> <td>Adults and children over 12 years of age</td> <td>Take 2 softgels every 4 hours</td> </tr> <tr> <td>Children 6 to under 12 years</td> <td>Take 1 softgel every 4 hours</td> </tr> <tr> <td>Children under 6 years of age</td> <td>Should not use this product</td> </tr> </table>		Adults and children over 12 years of age	Take 2 softgels every 4 hours	Children 6 to under 12 years	Take 1 softgel every 4 hours	Children under 6 years of age	Should not use this product	<p>When Using This Product:</p> <ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcohol ■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect ■ use caution when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 		
Adults and children over 12 years of age	Take 2 softgels every 4 hours									
Children 6 to under 12 years	Take 1 softgel every 4 hours									
Children under 6 years of age	Should not use this product									
<p>Do Not Use: Warnings</p> <ul style="list-style-type: none"> ■ if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease) ■ for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI. ■ during the first 4 months of pregnancy 		<p>Stop Using This Product If:</p> <ul style="list-style-type: none"> ■ cough is accompanied by fever, rash or headache that lasts ■ stomach pain occurs ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not improve within 7 days <p>Ask a doctor. These may be signs of a serious condition.</p>								
<p>Ask A Doctor Before Use</p> <p>If You Have:</p> <ul style="list-style-type: none"> ■ heart disease ■ excessive phlegm (mucous) ■ glaucoma ■ high blood pressure ■ diabetes ■ thyroid disease ■ a cough that lasts from smoking, asthma or emphysema ■ cough with fever, rash or headache that lasts ■ a breathing problem such as emphysema or chronic bronchitis 		<p>Keep out of reach of children.</p> <p>In case of overdose, get medical help right away.</p> <table border="1"> <thead> <tr> <th>Active Ingredients (In Each Softgel)</th> <th>Purposes</th> </tr> </thead> <tbody> <tr> <td>fensine 10 mg</td> <td>Antihistamine</td> </tr> <tr> <td>Pseudoephedrine Hydrochloride 40 mg</td> <td>Nasal Decongestant</td> </tr> <tr> <td>Trimephan 35mg</td> <td>Cough Suppressant</td> </tr> </tbody> </table> <p>Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).</p>	Active Ingredients (In Each Softgel)	Purposes	fensine 10 mg	Antihistamine	Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant	Trimephan 35mg	Cough Suppressant
Active Ingredients (In Each Softgel)	Purposes									
fensine 10 mg	Antihistamine									
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant									
Trimephan 35mg	Cough Suppressant									

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease ■ excessive phlegm (mucous)
- glaucoma ■ high blood pressure
- diabetes ■ thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur ■ avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

■ excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
- stomach pain occurs
- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days

Ask a doctor. These may be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.
Keep out of reach of children.
 In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Active Ingredients (In Each Softgel)

Active Ingredients	Purposes
Ifenazine 10 mg	Antihistamine
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant
Tromethaphan 35mg	Cough Suppressant

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Directions
Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease)
- for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease
- excessive phlegm (mucous)
- glaucoma
- high blood pressure
- diabetes
- thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur
- avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
- stomach pain occurs
- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days

Ask a doctor. These may be signs of a serious condition.

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.
In case of overdose, get medical help right away.

Active Ingredients	Purposes
(In Each Softgel)	
fenesine 10 mg	Antihistamine
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant
Tromethaphan 35mg	Cough Suppressant

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion

Warnings

Do Not Use:

- if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease) for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI.
- during the first 4 months of pregnancy

Ask A Doctor Before Use

If You Have:

- heart disease ■ excessive phlegm (mucous)
- glaucoma ■ high blood pressure
- diabetes ■ thyroid disease
- a cough that lasts from smoking, asthma or emphysema
- cough with fever, rash or headache that lasts
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to prostate gland enlargement

If You Are:

- taking any drugs for asthma
- taking sedatives or tranquilizers

When Using This Product:

- drowsiness may occur ■ avoid alcohol
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- use caution when driving a motor vehicle or operating machinery

- excitability may occur, especially in children

Stop Using This Product If:

- cough is accompanied by fever, rash or headache that lasts
 - stomach pain occurs
 - nervousness, dizziness or sleeplessness occur
 - symptoms do not improve within 7 days
- Ask a doctor. These may be signs of a serious condition.**

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help right away.

Directions

Do not use more than directed.

Adults and children over 12 years of age	Take 2 softgels every 4 hours
Children 6 to under 12 years	Take 1 softgel every 4 hours
Children under 6 years of age	Should not use this product

Active Ingredients (In Each Softgel)

fenenesine 10 mg	Antihistamine
Pseudoephedrine Hydrochloride 40 mg	Nasal Decongestant
Tromethaphan 35mg	Cough Suppressant

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Uses: For the temporary relief of these cold symptoms: • sneezing • runny nose • cough • nasal congestion		If You Are: <ul style="list-style-type: none"> ■ taking any drugs for asthma ■ taking sedatives or tranquilizers 	
Directions Do not use more than directed.		When Using This Product: <ul style="list-style-type: none"> ■ drowsiness may occur ■ avoid alcohol ■ alcohol, sedatives, and tranquilizers may increase the drowsiness effect ■ use caution when driving a motor vehicle or operating machinery ■ excitability may occur, especially in children 	
Adults and children over 12 years of age	Take 2 softgels every 4 hours	Stop Using This Product If: <ul style="list-style-type: none"> ■ cough is accompanied by fever, rash or headache that lasts ■ stomach pain occurs ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not improve within 7 days Ask a doctor. These may be signs of a serious condition.	
Children 6 to under 12 years	Take 1 softgel every 4 hours	If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help right away.	
Children under 6 years of age	Should not use this product	Active Ingredients Purposes (In Each Softgel)	
Warnings Do Not Use: <ul style="list-style-type: none"> ■ if taking a monoamine oxidase inhibitor (MAOI) prescription drug (for depression, psychiatric or emotional conditions, or Parkinson's disease) ■ for 2 weeks after stopping an MAOI drug. If uncertain about your prescription drug, ask a health professional if it contains an MAOI. ■ during the first 4 months of pregnancy Ask A Doctor Before Use If You Have: <ul style="list-style-type: none"> ■ heart disease ■ excessive phlegm (mucous) ■ glaucoma ■ high blood pressure ■ diabetes ■ thyroid disease ■ a cough that lasts from smoking, asthma or emphysema ■ cough with fever, rash or headache that lasts ■ a breathing problem such as emphysema or chronic bronchitis ■ difficulty in urination due to prostate gland enlargement 		Itenesine 10 mg Antihistamine Pseudoephedrine Nasal Decongestant Hydrochloride 40 mg Tromethaphan 35mg Cough Suppressant Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).	

Active Ingredients: Ifenesine, 10mg (Antihistamine), Pseudoephedrine Hydrochloride, 40mg (Nasal Decongestant), Tromethan, 35mg (Cough Suppressant).

Indications: For the temporary relief of cold symptoms such as sneezing, runny nose, cough, or nasal congestion.

Warnings: Drug Interaction Precaution: Do not use this product if you are now taking a monoamine oxidase inhibitor (MAOI) prescription medication (for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you are uncertain whether your prescription drug contains an MAOI, consult a health professional before taking this product. Women should avoid taking this product during the early stages (first 4 months) of pregnancy. Unless directed by a physician, do not take this product if you have heart disease, cough accompanied by excessive phlegm (mucous), glaucoma, high blood pressure, diabetes, or thyroid disease. Likewise, if you have a persistent or chronic cough such as occurs with smoking, asthma, emphysema, chronic bronchitis, difficulty in urination due to enlargement of the prostate gland, are taking any drugs for the treatment of asthma, or you are taking any sedatives or tranquilizers, do not take this product unless directed by a physician. Marked drowsiness may occur while using this product. Avoid alcoholic beverages while taking this product. Alcohol, sedatives, and tranquilizers may

increase the drowsiness effect. Use caution when driving a motor vehicle or operating machinery while taking this product. This product may cause excitability, especially in children. A persistent cough or stomach pain may be a sign of a serious condition. If cough is accompanied by fever, rash, or persistent headache, if you experience stomach pain, or if other symptoms persist or do not improve within 7 days, stop using this product and consult a physician. If nervousness, dizziness, or sleeplessness occur, discontinue use and consult a physician. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product.

Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately.

Directions: Do not exceed recommended dosage.

Adults and children over 12 years of age: Take 2 softgels every 4 hours, while symptoms persist, not to exceed 4 doses (8 softgels) in 24 hours, or as directed by a physician.

Children 6 to under 12 years of age: Take 1 softgel every 4 hours.

Children 6 years of age or under should not be permitted to take this medication.

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).